

Light tails and the Hermitian dual polar graphs

Jack Koolen*

School of Mathematical Sciences
University of Science and Technology of China,
Wen-Tsun Wu Key Laboratory of the Chinese Academy of Sciences,
230026, Anhui, PR China
e-mail: koolen@ustc.edu.cn

Zhi Qiao

School of Mathematical Sciences
University of Science and Technology of China,
230026, Anhui, PR China
email: gesec@mail.ustc.edu.cn

November 18, 2015

Dedicated to Andries Brouwer on the occasion of his 65th birthday

Abstract

Jurišić et al. [9] conjectured (see also [13, Problem 13]) that if a distance-regular graph Γ with diameter D at least three has a light tail, then one of the following holds:

1. $a_1 = 0$;
2. Γ is an antipodal cover of diameter three;
3. Γ is tight;
4. Γ is the halved $2D + 1$ -cube;
5. Γ is a Hermitian dual polar graph ${}^2A_{2D-1}(r)$ where r is a prime power.

In this note, we will consider the case when the light tail corresponds to the eigenvalue $-\frac{k}{a_1+1}$. Our main result is:

*JHK was partially supported by the National Natural Science Foundation of China (No. 11471009).

Theorem Let Γ be a non-bipartite distance-regular graph with valency $k \geq 3$, diameter $D \geq 3$ and distinct eigenvalues $\theta_0 > \theta_1 > \dots > \theta_D$. Suppose that Γ is 2-bounded with smallest eigenvalue $\theta_D = -\frac{k}{a_1+1}$. If the minimal idempotent E_D , corresponding to eigenvalue θ_D , is a light tail, then Γ is the dual polar graph ${}^2A_{2D-1}(r)$, where r is a prime power.

As a consequence of this result we will also show:

Theorem. Let Γ be a distance-regular graph with valency $k \geq 3$, diameter $D \geq 2$, $a_1 = 1$ and $\theta_0 > \theta_1 > \dots > \theta_D$. If $c_2 \geq 5$ and $\theta_D = -k/2$, then $c_2 = 5$ and Γ is the dual polar graph ${}^2A_{2D-1}(2)$.

1 Introduction

For definitions, see next section and [3]. Let Γ be a distance-regular graph with n vertices, valency $k \geq 3$ and diameter $D \geq 2$. A minimal idempotent E of Γ , say corresponding to eigenvalue θ , is called a light tail if $E \circ E = aE_0 + bF$, where $E_0 = \frac{1}{n}J$, $F \neq E_0$ is a minimal idempotent of Γ , and a, b are non-zero real numbers.

Jurišić et al. [9] conjectured (see also [13, Problem 13]) that if a distance-regular graph Γ with diameter D at least three has a light tail, then one of the following holds:

1. $a_1 = 0$;
2. Γ is an antipodal cover of diameter three;
3. Γ is tight;
4. Γ is the halved $2D + 1$ -cube;
5. Γ is a Hermitian dual polar graph ${}^2A_{2D-1}(r)$ where r is a prime power.

Jurišić et al. [9] showed that a distance-regular graph Γ with diameter at least three has a light tail corresponding to -1 if and only if Γ is an antipodal cover of diameter three. If a tight distance-regular graph has a light tail, then it is easy to show that Γ is antipodal (using the fact that tight implies that $a_D = 0$). In this note, we will consider distance-regular graphs with valency k and intersection number a_1 having a light tail corresponding to the eigenvalue $-\frac{k}{a_1+1}$. Our main result is:

Theorem 1.1. *Let Γ be a non-bipartite distance-regular graph with valency $k \geq 3$, diameter $D \geq 3$ and distinct eigenvalues $\theta_0 > \theta_1 > \dots > \theta_D$. Suppose that Γ is 2-bounded with smallest eigenvalue $\theta_D = -\frac{k}{a_1+1}$. If the minimal idempotent E_D , corresponding to eigenvalue θ_D , is a light tail, then Γ is the dual polar graph ${}^2A_{2D-1}(r)$, where r is a prime power.*

As a consequence of this result we will also show:

Theorem 1.2. *Let Γ be a distance-regular graph with valency $k \geq 3$, diameter $D \geq 2$, $a_1 = 1$ and distinct eigenvalues $\theta_0 > \theta_1 > \dots > \theta_D$. If $c_2 \geq 5$ and $\theta_D = -k/2$, then $c_2 = 5$ and Γ is the dual polar graph ${}^2A_{2D-1}(2)$.*

The authors [10] conjectured that if Γ is a distance-regular graph with valency $k \geq 3$, $a_1 = 1$, diameter $D \geq 3$ and smallest eigenvalue $-k/2$, then Γ is either the Hamming graph $H(D, 3)$, the dual polar graph $B_D(2)$ or the dual polar graph ${}^2A_{2D-1}(2)$ if D is large enough. We confirm this conjecture when $c_2 \geq 5$.

This note is organized as follows. In the next section, we give the definitions and some preliminary results. Then, in Section 3, we start with a sufficient condition for a distance-regular graph to have light tail, and then give a proof of Theorem 1.1. In the last section we will give a proof of Theorem 1.2.

2 Preliminaries and definitions

All graphs considered in this note are finite, undirected and simple (for more background information and undefined notions, see [3] or [13]). We denote $x \sim y$ if the vertices x, y are adjacent. For a connected graph $\Gamma = (V(\Gamma), E(\Gamma))$, the *distance* $d_\Gamma(x, y)$ between any two vertices x, y is the length of a shortest path between x and y in Γ , and the *diameter* $D(\Gamma)$ is the maximum distance between any two vertices of Γ , and if the graph is clear from the context, we use $d(x, y)$ and D for simplicity. For any vertex x , let $\Gamma_i(x)$ be the set of vertices in Γ at distance precisely i from x , where $0 \leq i \leq D$. A graph Γ is locally Δ , when for all vertices x the subgraph induced by $\Gamma_1(x)$ is isomorphic to Δ .

A connected graph Γ with diameter D is called a *distance-regular graph* if there are integers b_i, c_i ($0 \leq i \leq D$) such that for any two vertices $x, y \in V(\Gamma)$ with $d(x, y) = i$, there are exactly c_i neighbors of y in $\Gamma_{i-1}(x)$ and b_i neighbors of y in $\Gamma_{i+1}(x)$. The numbers b_i, c_i and $a_i := b_0 - b_i - c_i$ ($0 \leq i \leq D$) are called the *intersection numbers* of Γ . Set $c_0 = b_D = 0$. We observe $a_0 = 0$ and $c_1 = 1$. The array $\iota(\Gamma) = \{b_0, b_1, \dots, b_{D-1}; c_1, c_2, \dots, c_D\}$ is called the *intersection array* of Γ . In particular, Γ is a regular graph with valency $k := b_0$.

Let Γ be a distance-regular graph with n vertices, valency k and diameter D . For $0 \leq i \leq D$, let A_i be the $\{0, 1\}$ -matrix indexed by the vertices of Γ with $(A_i)_{xy} = 1$ if and only $d(x, y) = i$ for vertices x, y of Γ . We call A_i the *distance- i matrix* of Γ , and $A := A_1$ the *adjacency matrix* of Γ . The *eigenvalues* $\theta_0 > \theta_1 > \dots > \theta_D$ of Γ are just the eigenvalues of its adjacency matrix A . We denote m_i the *multiplicity* of θ_i .

For each eigenvalue θ_i of Γ , let W_i be a matrix with its columns forming an orthonormal basis for the eigenspace of θ . And $E_i := W_i W_i^T$ is called the *minimal idempotent* corresponding

to the eigenvalue θ_i , satisfying $E_i E_j = \delta_{ij} E_j$ and $A E_i = \theta_i E_i$, where δ_{ij} is the Kronecker symbol. Note that $n E_0$ is the all-one matrix J .

The set of distance matrices $\{A_0 = I, A_1, A_2, \dots, A_D\}$ forms a basis for a commutative $\mathbb{R}$ -algebra $\mathcal{A}$, known as the *Bose-Mesner algebra*. The set of minimal idempotents $\{E_0 = \frac{1}{n}J, E_1, E_2, \dots, E_D\}$ is another basis for $\mathcal{A}$. There exist $(D+1) \times (D+1)$ matrices P and Q (see [3, p.45]), such that the following relations hold:

$$A_i = \sum_{j=0}^D P_{ji} E_j \quad \text{and} \quad E_i = \frac{1}{n} \sum_{j=0}^D Q_{ji} A_j \quad (0 \leq i \leq D). \quad (1)$$

Note that $Q_{0i} = m_i$ (see [3, Lemma 2.2.1]). The Bose-Mesner algebra $\mathcal{A}$ is also closed under the Schur product, therefore we also have

$$E_i \circ E_j = \frac{1}{n} \sum_{h=0}^D q_{ij}^h E_h \quad (0 \leq i, j \leq D), \quad (2)$$

where the numbers q_{ij}^h ($0 \leq i, j, h \leq D$) are called the *Krein parameters* (see [3, p.48]). It is well known, cf. [3, Lemma 2.3.1, Theorem 2.3.2], the Krein parameters are non-negative real numbers and also $q_{ij}^0 = \delta_{ij} m_j$ holds.

Let $E_i = W_i W_i^T$ be a minimal idempotent corresponding to the eigenvalue θ_i , where the columns of W_i form an orthonormal basis of the eigenspace of θ_i . We denote the x -th row of W_i by $\hat{x}$. Note that $E_i \circ A_j = \frac{1}{n} Q_{ji} A_j$, hence all the vectors $\hat{x}$ has the same length $\frac{1}{n} Q_{0i}$ and the cosine of the angle between two vectors $\hat{x}$ and $\hat{y}$ is $u_j(\theta_i) := \frac{Q_{ji}}{Q_{0i}}$, where $d(x, y) = j$. The map $x \mapsto \hat{x}$ is called a *spherical representation* of Γ , and the sequence $\{u_j(\theta_i) \mid 0 \leq j \leq D\}$ is called the *standard sequence* of Γ , corresponding to the eigenvalue θ_i . As $AW = \theta_i E_i W = \theta_i W$, we have $\theta_i \hat{x} = \sum_{y \sim x} \hat{y}$, and hence the following holds:

$$\begin{aligned} c_j u_{j-1}(\theta_i) + a_j u_j(\theta_i) + b_j u_{j+1}(\theta_i) &= \theta_i u_j(\theta_i) \quad (1 \leq j \leq D-1) \\ c_D u_{D-1}(\theta_i) + a_D u_D(\theta_i) &= \theta_i u_D(\theta_i) \end{aligned} \quad (3)$$

with $u_0(\theta_i) = 1$ and $u_1(\theta_i) = \frac{\theta_i}{k}$.

A minimal idempotent E is called a *light tail* if $E \circ E = aE_0 + bF$ for some minimal idempotent $F \neq E_0$ and nonzero real numbers a, b . The minimal idempotent F is called the *associated minimal idempotent* to the light tail E . Note that $E_0 \circ E_0 = \frac{1}{n} E_0$ holds and that if Γ is bipartite, then also $E_D \circ E_D = \frac{1}{n} E_0$ holds, see for example [11, Lemma 3.3]. So the eigenvalue corresponding to the light tail E is different from $\pm k$.

Jurišić et al. [9, Theorem 3.2] has shown the following result.

Lemma 2.1. *Let Γ be a distance-regular graph with valency $k \geq 3$, diameter $D \geq 2$ and distinct eigenvalues $\theta_0 > \theta_1 > \dots > \theta_D$. Let $\theta_i \neq \pm k$ be an eigenvalue of Γ , then*

$$\frac{m_i - k}{k} \geq -\frac{(\theta_i + 1)^2 a_1 (a_1 + 1)}{((a_1 + 1)\theta_i + k)^2 + k a_1 b_1}, \quad (4)$$

and equality holds if and only if E_i is a light tail.

For a graph Γ , a partition $\Pi = \{P_1, P_2, \dots, P_t\}$ of $V(\Gamma)$ is called *equitable* if there are constants α_{ij} ($1 \leq i, j \leq t$) such that all vertices $x \in P_i$ have exactly α_{ij} neighbours in P_j . The α_{ij} 's ($1 \leq i, j \leq t$) are called the parameters of the equitable partition.

Let Γ be a distance-regular graph. For a set S of vertices of Γ , define $S_i = \{x \in V(\Gamma) \mid d(x, S) = i\}$, where $d(x, S) := \min\{d(x, y) \mid y \in S\}$. Then number $\rho(S) := \max\{i \mid S_i \neq \emptyset\}$ is called the *covering radius* of S . The set S is called a *completely regular code* of Γ if the distance-partition $\{S = S_0, S_1, S_2, \dots, S_{\rho(S)}\}$ is equitable.

The following result was first shown by Delsarte [4] for strongly regular graphs, and extended by Godsil [5] to distance regular graphs.

Lemma 2.2. *Let Γ be a distance-regular graph with valency k and smallest eigenvalue $\theta_{\min}$. Let C be a clique in Γ with c vertices. Then $c \leq 1 - \frac{k}{\theta_{\min}}$, and equality holds if and only if C is a completely regular code with covering radius $D - 1$.*

A clique C in a distance-regular graph Γ that attains the above bound is called a *Delsarte clique*. A distance-regular graph Γ is called *geometric* (with respect to $\mathcal{C}$) if it contains a collection $\mathcal{C}$ of Delsarte cliques such that each edge is contained in a unique $C \in \mathcal{C}$.

Let Γ be a geometric distance-regular graph with respect to $\mathcal{C}$. As each edge is contained in a unique Delsarte clique in $\mathcal{C}$, we see the Delsarte cliques have size $a_1 + 2$ and $\theta_{\min} = -\frac{k}{a_1 + 1}$, where $\theta_{\min}$ is the smallest eigenvalue of Γ . And the parameter $\gamma_i := |\Gamma_i(x) \cap C|$ ($0 \leq i \leq D$) is well-defined (see [5, Theorem 7.1]), where C is any Delsarte clique of Γ and x is any vertex with $d(x, C) = i$.

Lemma 2.3. (cf. [13, Proposition 4.3]) *Let Γ be a geometric distance-regular graph with valency $k \geq 3$, diameter $D \geq 2$ and distinct eigenvalues $\theta_0 > \theta_1 > \dots > \theta_D$. Then the following holds:*

$$\gamma_i u_i(\theta_D) + (a_1 + 2 - \gamma_i) u_{i+1}(\theta_D) = 0 \quad (0 \leq i \leq D - 1). \quad (5)$$

where $u_i(\theta_D)$ ($0 \leq i \leq D$) is the standard sequence corresponding to θ_D .

A subgraph Δ of a graph Γ is called *strongly closed* if, for all vertices $x, y \in V(\Delta)$ and $z \in V(\Gamma)$ such that $d_\Gamma(x, z) + d_\Gamma(z, y) \leq d_\Gamma(x, y) + 1$, holds, then we have that $z \in V(\Delta)$. A graph Γ is said to be *m-bounded* for some $1 \leq m \leq D$, if for all $1 \leq i \leq m$ and all vertices x, y at distance i , there exists a strongly closed subgraph $\Delta(x, y)$ of Γ with diameter i , containing x and y as vertices.

The following result was shown by Hiraki [7, Theorem 1.1], see also [13, Proposition 11.3].

Lemma 2.4. *Let Γ be a distance-regular graph with diameter $D \geq 3$. Let $m \in \{1, 2, \dots, D-1\}$ and $c_{m+1} \neq 1$. Then Γ is m -bounded if and only if Γ is $K_{1,1,2}$ -free, $a_1 \neq 0$, $a_i = c_i a_1$ holds for $i = 1, 2, \dots, m$ and $c_{m-1} < c_m$.*

A generalized quadrangle of order (s, t) , denoted by $\text{GQ}(s, t)$, (where s, t are positive integers), is a $K_{2,1,1}$ -free distance-regular graph with diameter 2 with intersection array $\{(t+1)s, ts; 1, t+1\}$.

3 Characterization of the Hermitian dual polar graphs

We will give a characterization of the Hermitian dual polar graphs ${}^2A_{2D-1}(r)$, where r is a prime power. First we give a sufficient condition for a minimal idempotent to be a light tail of a distance-regular graph.

Lemma 3.1. *Let Γ be a distance-regular graph with n vertices, valency $k \geq 3$, diameter $D \geq 2$, $a_1 \neq 0$ and distinct eigenvalues $\theta_0 > \theta_1 > \dots > \theta_D = -\frac{k}{a_1+1}$. Then $\theta_1 \geq \frac{k-(a_1+1)(a_1+2)}{(a_1+1)^2}$ and equality holds if and only if E_D is a light tail with associated minimal idempotent E_1 .*

Proof. For each eigenvalue θ_i of Γ , we denote m_i its multiplicity and $u_j(\theta_i)$ ($0 \leq j \leq D$) its standard sequence.

By the definition of standard sequence, we have

$$\frac{n}{m_i} E_i = \sum_{j=0}^D u_j(\theta_i) A_j \quad (0 \leq i \leq D). \quad (6)$$

As $A_i \circ A_j = \delta_{ij} A_j$, the following holds

$$\left(\frac{n}{m_D} E_D\right) \circ \left(\frac{n}{m_D} E_D\right) = \sum_{i=0}^D u_i(\theta_D)^2 A_i. \quad (7)$$

By (2) and (6), we see

$$\left(\frac{n}{m_D} E_D\right) \circ \left(\frac{n}{m_D} E_D\right) = \frac{n^2}{m_D^2} \cdot \frac{1}{n} \sum_{i=0}^D q_{DD}^i E_i \quad (8)$$

$$= \frac{n}{m_D} E_0 + \frac{n}{m_D^2} \sum_{i=1}^D q_{DD}^i E_i \quad (9)$$

$$= \frac{1}{m_D} J + \sum_{j=0}^D \left(\sum_{i=1}^D \frac{m_i q_{DD}^i}{m_D^2} u_j(\theta_i) \right) \cdot A_j \quad (10)$$

$$= \frac{1}{m_D} J + \frac{m_D - 1}{m_D} \tilde{E}, \quad (11)$$

where $\tilde{E} = \sum_{j=0}^D \tilde{u}_j A_j$, $\tilde{u}_j = \sum_{i=1}^D \alpha_i u_j(\theta_i)$ and $\alpha_i = \frac{m_i q_{DD}^i}{m_D(m_D-1)} \geq 0$.

Since $J = \sum_{i=0}^D A_i$, by (7) and (11), we have

$$u_i^2(\theta_D) = \frac{1}{m_D} + \frac{m_D - 1}{m_D} \tilde{u}_i \quad (0 \leq i \leq D). \quad (12)$$

Since $u_0(\theta_D) = 1$ and $u_1(\theta_D) = \frac{\theta_D}{k} = -\frac{1}{a_1+1}$, we see that $\tilde{u}_0 = 1$ and $\tilde{u}_1 = \frac{m_D(\frac{1}{a_1+1})^2 - 1}{m_D - 1}$. As $a_1 \neq 0$, we see $\theta_D = -\frac{k}{a_1+1} \neq -k$. And Lemma 2.1 implies $m_D \geq \frac{a_1 k}{a_1+1} + 1$. Hence we have $\tilde{u}_1 \geq \frac{k-(a_1+1)(a_1+2)}{(a_1+1)^2 k}$. Note that $1 = \tilde{u}_0 = \sum_{i=1}^D \alpha_i$ and $\tilde{u}_1 = \sum_{i=1}^D \alpha_i u_1(\theta_i) = \frac{1}{k} \sum_{i=1}^D \alpha_i \theta_i$. We have that $\theta_1 \geq k \tilde{u}_1 \geq \frac{k-(a_1+1)(a_1+2)}{(a_1+1)^2}$, and equality holds if and only if $\tilde{E} = E_1$ and

$$\left(\frac{n}{m_D} E_D\right) \circ \left(\frac{n}{m_D} E_D\right) = \frac{1}{m_D} J + \frac{m_D - 1}{m_D} E_1,$$

both hold, by (11). This means that equality holds if and only if E_D is a light tail with associated minimal idempotent E_1 , by the definition of a light tail. $\square$

The following result is a consequence of [1, Cor 3.4].

Lemma 3.2. *Let Γ be a distance-regular graph with valency $k \geq 3$, diameter $D \geq 2$ and distinct eigenvalues $\theta_0 > \theta_1 > \dots > \theta_D$. If $c_2 \neq 1$ and Γ contains an induced $GQ(s, c_2 - 1)$ for some positive integer s , then $\theta_1 \leq \frac{k-(a_1+1)}{c_2-1} - 1$.*

Proof. As Γ contains an induced subgraph $\Gamma' = GQ(s, c_2 - 1)$, then, by [1, Corollary 3.4], we see $-c_2 = \theta_{\min}(\Gamma') \geq -1 - \frac{b_1}{\theta_1+1}$. The result follows. $\square$

The following result follows immediately from Lemmata 3.1 and 3.2.

Theorem 3.3. *Let Γ be a distance-regular graph with valency $k \geq 3$, diameter $D \geq 2$, $a_1 \neq 0$ and distinct eigenvalues $\theta_0 > \theta_1 > \dots > \theta_D = -\frac{k}{a_1+1}$. If Γ contains an induced $GQ(a_1 + 1, c_2 - 1)$ with $c_2 = (a_1 + 1)^2 + 1$, then $\theta_1 = \frac{k-(a_1+1)(a_1+2)}{(a_1+1)^2}$ and E_D is a light tail with associated minimal idempotent E_1 .*

Proof. Since $c_2 = (a_1 + 1)^2 + 1 \neq 1$ and Γ contains an induced $GQ(a_1 + 1, c_2 - 1)$, we have $\theta_1 \leq \frac{k-(a_1+1)}{c_2-1} - 1 = \frac{k-(a_1+1)}{(a_1+1)^2} - 1$, by Lemma 3.2. Lemma 3.1 implies $\theta_1 \geq \frac{k-(a_1+1)}{(a_1+1)^2} - 1$. These two bounds on θ_1 imply that $\theta_1 = \frac{k-(a_1+1)}{(a_1+1)^2} - 1$. Now again by Lemma 3.1 it follows that E_D is a light tail with associated minimal idempotent E_1 . $\square$

Proof of Theorem 1.1. Since E_D is a light tail with $\theta_D = -\frac{k}{a_1+1}$, the graph Γ is locally the disjoint union of $(a_1 + 1)$ -cliques by [9, Corollary 6.3]. This implies that all maximal

cliques of Γ are $(a_1 + 2)$ -cliques, and hence Delsarte cliques by Lemma 2.2. It follows that Γ is geometric.

Note that we have the following equations:

$$c_i + a_i + b_i = k, \quad (0 \leq i \leq D) \quad (13)$$

$$c_i u_{i-1}(\theta_D) + a_i u_i(\theta_D) + b_i u_{i+1}(\theta_D) = \theta_D u_i(\theta_D), \quad (1 \leq i \leq D-1) \quad (14)$$

$$\gamma_i u_i(\theta_D) + (a_1 + 2 - \gamma_i) u_{i+1}(\theta_D) = 0, \quad (0 \leq i \leq D-1) \quad (15)$$

where Equation (15) is from Lemma 2.3.

As $u_i(\theta_D) \neq 0$ for $0 \leq i \leq D$ by [3, Corollary 4.1.2] we see that $\gamma_i \leq a_1 + 1$ for $0 \leq i \leq D-1$. Then Equations (13) (14) and (15) implies the following:

$$a_i = c_i \frac{a_1 + 1 - \gamma_{i-1}}{\gamma_{i-1}} + b_i \frac{\gamma_i - 1}{a_1 + 1 - (\gamma_i - 1)} \quad (1 \leq i \leq D-1) \quad (16)$$

$$= c_i \frac{a_1 + 1 - \gamma_{i-1} - (\gamma_i - 1)}{\gamma_{i-1}} + k \frac{\gamma_i - 1}{a_1 + 1} \quad (1 \leq i \leq D-1) \quad (17)$$

Now we assume Γ is m -bounded for some $2 \leq m \leq D-2$. We will show that Γ is $(m+1)$ -bounded, and this implies, by induction, that Γ is $(D-1)$ -bounded.

We claim that $\gamma_i = 1$ ($i \leq m$) and $a_i = c_i a_1$ ($i \leq m$).

First, we consider the case $c_{m+1} = 1$. Choose a vertex x and a Delsarte clique C with $d(x, C) = i$. As $\gamma_i \leq a_1 + 1$, we can choose a vertex $y \in C$ with $d(x, y) = i + 1$. We find $C \cap \Gamma_i(x) \subseteq \Gamma(y) \cap \Gamma_i(x)$, i.e. $c_{i+1} \geq \gamma_i$ ($i \leq m$). Note that $c_1 \leq c_2 \leq \dots \leq c_D$ by [3, Proposition 4.1.6]. We have $\gamma_i = 1$ ($i \leq m$), which implies $a_i = c_i a_1$ by Equation (16).

In the case $c_{m+1} \neq 1$, as Γ is m -bounded, by Lemma 2.4, we have $a_i = c_i a_1$ ($i \leq m$). Note that $\gamma_0 = 1$. Assume $\gamma_{i-1} = 1$ ($1 \leq i \leq m$), with $a_i = c_i a_1$, we have $\gamma_i = 1$. By induction, it follows that $\gamma_i = 1$ ($i \leq m$).

As E_D is a light tail with associated minimal idempotent F corresponding to the eigenvalue $\theta' \neq k$ and $\theta_D = -\frac{k}{a_1+1}$, by [9, Theorem 4.1], we have

$$u_i(\theta_D)^2 = \alpha + \beta u_i(\theta') \quad (0 \leq i \leq D), \quad (18)$$

where $\theta' = \frac{k-(a_1+1)(a_1+2)}{(a_1+1)^2}$, $\alpha = \frac{a_1+1}{a_1 k + a_1 + 1}$ and $\beta = \frac{a_1 k}{a_1 k + a_1 + 1}$.

For $i \leq m \leq D-2$, we have

$$u_{i+1}(\theta_D) = \left(-\frac{1}{a_1 + 1}\right)^{i+1} \quad (0 \leq i \leq m), \quad (19)$$

$$u_{i+1}(\theta') = \frac{(a_1 + 1)^{-2(i+1)}(a_1 k + a_1 + 1) - (a_1 + 1)}{k a_1} \quad (0 \leq i \leq m), \quad (20)$$

by Equations (15) and (18), as $\gamma_i = 1$. Substitute $a_i = c_i a_1$, $b_i = k - c_i(a_1 + 1)$ ($1 \leq i \leq m$) and Equation (20) into

$$c_i u_{i-1}(\theta') + a_i u_i(\theta') + b_i u_{i+1}(\theta') = \theta' u_i(\theta') \quad (1 \leq i \leq D), \quad (21)$$

to obtain

$$c_i = \frac{(a_1 + 1)^{2i} - 1}{(a_1 + 1)^2 - 1} \quad (0 \leq i \leq m). \quad (22)$$

It follows that $c_2 = (a_1 + 1)^2 + 1 \neq 1$. Since Γ is m -bounded with $m \geq 2$ and $c_2 \neq 1$, [8, Proposition 19] implies $c_{m+1} \geq c_m + (c_m - c_{m-1})(c_2 - c_1) > c_m$, i.e.

$$c_{m+1} \geq \frac{(a_1 + 1)^{2(m+1)} - 1}{(a_1 + 1)^2 - 1}. \quad (23)$$

Equations (15) and (18) imply

$$u_{m+2}(\theta_D) = \left(-\frac{\gamma_{m+1}}{a_1 + 2 - \gamma_{m+1}}\right) u_{m+1}(\theta_D), \quad (24)$$

$$u_{m+2}(\theta') = \frac{\gamma_{m+1}^2 (a_1 + 2 - \gamma_{m+1})^{-2} (a_1 + 1)^{-2m-2} (a_1 k + a_1 + 1) - (a_1 + 1)}{k a_1}. \quad (25)$$

As Γ is 2-bounded and $c_2 \neq 1$, Lemma 2.4 implies that Γ contains an induced $GQ(a_1 + 1, c_2 - 1)$. As Γ is not bipartite, we see $\theta_D \neq -k$ by [6, Theorem 8.8.2]. And $\theta_D = -\frac{k}{a_1 + 1}$ implies $a_1 \neq 0$. It follows that $\theta' = \theta_1$ from Theorem 3.3. Then we have $u_0(\theta') > u_1(\theta') > \dots > u_D(\theta')$ by [5, Theorem 13.2.1]. From Equations (18) and (24), we see

$$\gamma_{m+1} \leq \frac{a_1 + 1}{2}, \quad (26)$$

as $u_{m+1}(\theta') > u_{m+2}(\theta')$.

Substitute Equations (17), (20) and (25) into (21), we obtain

$$c_{m+1} = \frac{(a_1 + 2 - \gamma_{m+1})(a_1 + 1)((a_1 + 1)^{2(m+1)} - 1) - k(a_1 + 2)(\gamma_{m+1} - 1)}{(a_1 + 1)^2(a_1 + 2)(a_1 + 1 - \gamma_{m+1})}. \quad (27)$$

Substituting Equation (27) into (23), we see that

$$\frac{(\gamma_{m+1} - 1)((a_1 + 1)((a_1 + 1)^{2m+2} - 1) - ((a_1 + 1)^2 - 1)k)}{a_1(a_1 + 1)^2(a_1 + 2)(a_1 + 1 - \gamma_{m+1})} \geq 0. \quad (28)$$

As $\gamma_m = 1$, by Equation (16), we see that $a_{m+1} \geq a_1 c_{m+1}$. Since $m \leq D - 2$, we see that $k > a_{m+1} + c_{m+1} \geq (a_1 + 1)c_{m+1}$ and Equation (23) implies

$$k > (a_1 + 1) \frac{(a_1 + 1)^{2(m+1)} - 1}{(a_1 + 1)^2 - 1}. \quad (29)$$

We see $\gamma_{m+1} = 1$ by Equations (26) (29) and (28). It follows that $c_{m+1} = \frac{(a_1+1)^{2(m+1)}-1}{(a_1+1)^2-1} > c_m$ by Equations (27) and (22), and $a_{m+1} = c_{m+1}a_1$ by Equation (17). This shows that Γ is $(m+1)$ -bounded by Lemma 2.4.

Now we have that Γ is $(D-1)$ -bounded. Note that $\gamma_{D-1} = 1$. By the same way to obtain Equations (16), (17), and (20), we have

$$a_D = (a_1 + 1)c_D - \frac{k}{a_1 + 1} \quad (30)$$

$$= a_1 c_D, \quad (31)$$

$$u_D(\theta') = \frac{(a_1 + 1)^{-2D}(a_1 k + a_1 + 1) - (a_1 + 1)}{k a_1}. \quad (32)$$

Substitute $k = (a_1 + 1)c_D$, $a_D = c_D a_1$ and Equations (20) (32) into

$$c_D u_{D-1}(\theta') + a_D u_D(\theta') = \theta' u_D(\theta'), \quad (33)$$

to obtain that

$$c_D = \frac{(a_1 + 1)^{2D} - 1}{(a_1 + 1)^2 - 1} \quad (34)$$

holds. Now we see that Γ is a distance-regular graph with $a_i = c_i a_1$ and $c_i = \frac{(a_1+1)^{2i}-1}{(a_1+1)^2-1}$ ($i \leq D$). As Γ has diameter $D \geq 3$, by [3, Theorem 9.4.7], Γ is ${}^2A_{2D-1}(r)$ with $r = a_1 + 1$. $\square$

Remark. The Hermitian dual polar graph ${}^2A_{2D-1}(r)$ is determined by its intersection array for diameter $D \geq 3$ (see [3, Theorem 9.4.7]). The graph ${}^2A_3(r)$ is a generalized quadrangle $GQ(r, r^2)$. It is known that for r a prime power with $r \equiv 2 \pmod{3}$ and $r > 2$, there are at least two non-isomorphic generalized quadrangles $GQ(r, r^2)$'s, so ${}^2A_3(r)$ is not determined by its intersection array for those prime powers (see [12, 3.1.2, 3.1.6 and 3.2.5]).

4 Proofs of Theorem 1.2

First we establish the following consequence of Theorem 1.1.

Corollary 4.1. *Let Γ be a non-bipartite geometric distance-regular graph with valency $k \geq 3$, diameter $D \geq 3$ and eigenvalues $\theta_0 > \theta_D > \dots > \theta_D$. If Γ is m -bounded for some $m \geq 2$ and $c_2 \geq (a_1 + 1)^2 + 1$, then $c_2 = (a_1 + 1)^2 + 1$ and Γ is the dual polar graph ${}^2A_{2D-1}(r)$ with $r = a_1 + 1$.*

Proof. The graph Γ is non-bipartite geometric implies that $\theta_D = -\frac{k}{a_1+1}$ and $a_1 \neq 0$. As Γ is 2-bounded with $c_2 \geq (a_1 + 1)^2 + 1 \neq 1$, by Lemma 2.4, we see that Γ contains an induced

subgraph $GQ(a_1 + 1, c_2 - 1)$ and hence $\theta_1 \leq \frac{k-(a_1+1)}{c_2-1} - 1$, by Lemma 3.2. Lemma 3.1 implies that $c_2 = (a_1 + 1)^2 + 1$ and now it follows that E_D is a light tail with associated minimal idempotent E_1 , by Theorem 3.3. Now, the result follows from Theorem 1.1. $\square$

Proof of Theorem 1.2. Every triangle of Γ is a Delsarte clique and Γ is geometric.

Choose a vertex x and a Delsarte clique C with $d(x, C) = i$. Then there exists a vertex $y \in \Gamma(x)$ such that $d(y, C) = i - 1$. Note that $\Gamma_{i-1}(y) \cap C \subseteq \Gamma_i(x) \cap C$, which implies $\gamma_i \geq \gamma_{i-1}$ ($1 \leq i \leq D$). So the sequence γ_i is non-decreasing and $1 \leq \gamma_i \leq a_1 + 1 = 2$.

Then there exists some integer e such that

$$\gamma_i = \begin{cases} 1, & i \leq e - 1, \\ 2, & i \geq e. \end{cases} \quad (35)$$

And (5) implies the following

$$u_i(\theta_D) = \begin{cases} (-\frac{1}{2})^i, & i \leq e, \\ (-\frac{1}{2})^{2e-i}, & i \geq e + 1. \end{cases} \quad (36)$$

As $|u_i(\theta_D)| \leq 1$, which implies $e \geq \frac{D}{2}$. If $D \geq 5$, then $e \geq 3$. Then $a_i = c_i a_1$ ($i \leq e - 1$) by Equation (16). Lemma 2.4 implies that Γ is 2-bounded, hence it follows that $c_2 = 5$ and Γ is the dual polar graph ${}^2A_{2D-1}(2)$ by Corollary 4.1. The case when $2 \leq D \leq 4$ follows from [10, Theorem 1.2]. $\square$

Acknowledgements. We would like to thank Jongyook Park. He carefully read an earlier version of the note and his comments improved the note significantly.

References

- [1] S. Bang, J.H. Koolen, and J. Park. Some results on the eigenvalues of distance-regular graphs. *Graphs and Combin.*, 2015.
- [2] A.E. Brouwer. Parameters of strongly regular graphs. <http://www.win.tue.nl/~aeb/graphs/srg/srgtab.html>, September 2015.
- [3] A.E. Brouwer, A.M. Cohen, and A. Neumaier. *Distance-Regular Graphs*. Springer-Verlag, Berlin, 1989.
- [4] P. Delsarte. *An algebraic approach to the association schemes of coding theory*, volume 10 of *Philips Res. Reports Suppl.* Philips, 1973.
- [5] C.D. Godsil. *Algebraic Combinatorics*. Chapman & Hall, New York, 1993.

- [6] C.D. Godsil and G. Royle. *Algebraic Graph Theory*, volume 207 of *GTM*. Springer, New York, 2001.
- [7] A. Hiraki. Strongly closed subgraphs in a regular thick near polygon. *European J. Combin.*, 20:789–796, 1999.
- [8] A. Hiraki. A characterization of some distance-regular graphs by strongly closed subgraphs. *European J. Combin.*, 30:893–907, 2009.
- [9] A. Jurišić, P. Terwilliger, and A. Žitink. Distance-regular graphs with light tails. *European J. Combin.*, 31:1539–1552, 2010.
- [10] J.H. Koolen and Z. Qiao. Distance-regular graph with valency k and smallest eigenvalue at most $-\frac{k}{2}$. [arXiv:1507.04839v2].
- [11] A.A. Pascasio. Tight graphs and their primitive idempotents. *J. Algebraic Combin.*, 10:47–59, 1999.
- [12] S.E. Payne and J.A. Thas. *Finite Generalized Quadrangles*. European Mathematical Society, second edition, 2009.
- [13] E.R. van Dam, J.H. Koolen, and H. Tanaka. Distance-regular graphs. [arXiv:1410.6294v1].